










COVID-19 Vaccines

	PFIZER	MODERNA	JOHNSON & JOHNSON
 AGE	16 years and older	18 years and older	18 years and older
 DOSAGE	2 doses, 21 days apart	2 doses, 28 days apart and older	1 dose
 COMMON SIDE EFFECTS	Injection site pain, fatigue, headache, muscle pain, joint pain, and fever	Injection site pain, fatigue, headache, muscle pain, joint pain, and fever	Injection site reactions, headache, fatigue, myalgia, nausea, and fever
 HOW EFFECTIVE	According to clinical trial data, all 3 authorized vaccines are 100% effective at preventing hospitalization and death related to COVID-19.		
 SIDE EFFECTS ARE NORMAL	Side effects are normal and are a sign that your body is developing immunity against the virus that causes COVID-19. They should go away within a day or two. Not everyone will experience side effects.		
 CHRONIC CONDITIONS	If you have chronic health conditions you can still get this vaccine. You are more at risk of having complications from COVID-19 if you get infected with the virus.		
 PREGNANCY	Women who are pregnant, breastfeeding or are trying to become pregnant are also advised to receive the vaccine. Getting infected with COVID-19 while pregnant can cause medical complications to the mother and baby. Please talk to your doctor about getting the vaccine if you have more questions.		

Due to limited COVID-19 vaccines, individuals are encouraged to receive the vaccine that is available at the time. The 3 authorized vaccines have been proven to be safe. *Source: Centers for Disease Control and Prevention (2021)